AD-A102 427   NEW YORK UNIV N Y DEPT OF CHEMISTRY   F/6 7/3
              PHOTOELECTRON EMISSION SPECTROSCOPY OF WEAK ACIDS AND BASES AND--ETC(U)
              JUN 81 K VON BURG, P. DELAHAY   N00014-75-C-0397
UNCLASSIFIED TR-17   NL

END
DATE
FILMED
8 – 81
DTIC

AD A102427

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0397

Task No. NR 051-258

TECHNICAL REPORT NO. 17 (New Series)

Photoelectron Emission Spectroscopy of

Weak Acids and Bases and Their Ions in Aqueous Solution,

by

Kathrin/von Burg and Paul/Delahay

Prepared for Publication

in

Chemical Physics Letters

New York University
Department of Chemistry
New York, NY

AUG 4 1981

A

June 1981

81 8 04 024

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER 17 (New Series)	2. GOVT ACCESSION NO. AD-A102427	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) PHOTOELECTRON EMISSION SPECTROSCOPY OF WEAK ACIDS AND BASES AND THEIR IONS IN AQUEOUS SOLUTION	5. TYPE OF REPORT & PERIOD COVERED Technical Report	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) Kathrin von Burg and Paul Delahay	8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0397	
9. PERFORMING ORGANIZATION NAME AND ADDRESS New York University New York, NY 10003	10. PROGRAM ELEMENT. PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-258	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217	12. REPORT DATE June 1981	
	13. NUMBER OF PAGES 22	
14. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office)	15. SECURITY CLASS. (of this report)	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	

16. DISTRIBUTION STATEMENT (of this Report)

This document has been approved for public release and sale; its distribution is unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Acid, weak   Emission
Aqueous solution   Photoelectron spectroscopy
Base, weak
Electron transfer

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

Threshold energies E_t are determined for 14 weak acids and three bases and their ions. Values of E_t are interpreted in terms of dissociation or protonation reactions involving species produced by photoionization. Electron transfer to weak acid molecules in solution is shown to occur for emission by inorganic anions and cations and weak acids.

PHOTOELECTRON EMISSION SPECTROSCOPY OF WEAK ACIDS AND BASES AND THEIR IONS IN
AQUEOUS SOLUTION

KATHRIN von BURG and PAUL DELAHAY

Department of Chemistry, New York University, New York, New York 10003, U.S.A.

Threshold energies E_t are determined for 14 weak acids and three bases
and their ions. Values of E_t are interpreted in terms of dissociation or
protonation reactions involving species produced by photoionization. Electron
transfer to weak acid molecules in solution is shown to occur for emission by
inorganic anions and cations and weak acids.

The threshold energies E_t for photoelectron emission by aqueous
solutions of a few anions of weak acids were reported in earlier papers [1,2].
No correlation of the E_t values was attempted at that stage. This is done
in the present paper on the basis of rather extensive data for weak acids and
bases and their ions. Weak acids will also be shown to be excellent electron
acceptors in solution.

1. Weak acids and their anions

The experimental methods developed in [1] and applied in [2-4] were used
in the present work. Emission spectra obtained for aqueous solutions of acids
and their salts displayed the yield Y (collected electrons per incident
photon) as a function of the photon energy E. The threshold energies E_t^j
(table 1) were obtained by extrapolation of the linear plots Y^n vs. E to Y =
0 for n = 0.5 or 0.4 [5]. The choice of exponent is discussed in [2-5]. The
yield for formic acid (4 M) was very low and only about twice that for pure
water (E_t = 10.06 eV [4]).

The threshold energies will be interpreted on the basis of the equation
[2-4],

$$\Delta G_e^j = \Delta G_H + \Delta G^j - \Delta G_r^j, \tag{1}$$

for the free energy of emission ΔG_e^j of the species $H_{n-j}A^{j-}$ (aq) (n = 1 to 3, j = 0 to 3). The quantity $\Delta G_H = 4.50$ eV [6] is the free energy of formation of the electron in vacuum on the assumption that the surface potential at the solution-water vapor interface is negligible ($\pm$ 0.1 V for water [7]). ΔG^j is the change of free energy for the reaction

$$H_{n-j}A^{j-}(aq) + H^+(aq) = H_{n-j}A^{(j-1)-}(aq) + 1/2H_2(g), \tag{2}$$

and ΔG_r^j (< 0) is the reorganization free energy for the process in which there is vibrational relaxation and the solvation nuclear configuration adjusts from j- to (j-1)- ionic charge.

Equation (1) will be applied first to the acid H_nA and its anion $H_{n-1}A^-$. One has (fig 1)

$$\Delta G_e(H_nA) - \Delta G_e(H_{n-1}A^-) = [\Delta G(H_{n-1}A^-) - \Delta G(H_nA)]$$
$$+ [\Delta G(H_nA^+) - \Delta G(H_{n-1}A)] + [\Delta G_r(H_{n-1}A) - \Delta G_r(H_nA^+)] \tag{3}$$

where the symbol $\Delta G(X)$ represents the free energy of formation of species X in aqueous solution, and the ΔG_r's are the reorganization free energies. Equation (3) is similar to the corresponding relationship for water and hydroxide ion [4]. Each of the three terms on the right hand side of eq 3 will be discussed.

One has

$$\Delta G_K^1 = \Delta G(H_{n-1}A^-) - \Delta G(H_nA)$$
$$= - 0.0592 \log K_1 \quad , \tag{4}$$

where ΔG_K^1 is expressed in electronvolts and K_1 is the first dissociation constant of H_nA. One has $0.1 < \Delta G_K^1 < 0.3$ eV for the acids of table 1. The third terms between brackets on the right hand side in eq (3) pertains to the change of ionic charge from -1 to 0 and 0 to 1 in the

photoionization of $H_{n-1}A^-$ and H_nA, respectively. The ions $H_{n-1}A^-$ and H_nA^+ have comparable sizes although the latter is presumably slightly bulkier than the former. Contributions from vibrational relaxation should nearly cancel out in the difference of ΔG_r's. Since the ΔG_r's are negative, one concludes on the basis of ionic sizes and equality of charge in absolute value that the third term in eq (3) is negative and small (a few tenths of electronvolt).

Equation (3) can be rewritten in the form,

$$\Delta^1 = E_t^o - E_t^1 - \Delta G_K^1$$
$$= [\Delta G(H_nA^+) - \Delta G(H_{n-1}A)] + \Delta^1(\Delta G_r), \tag{5}$$

where the free energies of emission are equated to the threshold energies [2-4] and the difference of reorganization free energies is denoted by $\Delta^1(\Delta G_r)$. The superscripts o and 1 correspond to H_nA and $H_{n-1}A^-$, respectively. The first term between brackets in eq (5) is the change of free energy for the protonation reaction,

$$H_{n-1}A + H^+ = H_nA^+, \tag{6}$$

involving the species produced by photoionization of $H_{n-1}A^-$ and H_nA, respectively. The ion H_nA^+ is thermodynamically unstable and the radical $H_{n-1}A$ does not protonate to any significant extent if the condition, $\Delta^1 - \Delta^1(\Delta G_r) > 0$, is satisfied (eq (5)). Since $\Delta^1(\Delta G_r)$ is negative and $\Delta^1 > -0.2$ eV, the foregoing conclusion is fulfilled for all the acids of table 1, albeit marginally for maleic acid. Thus, <u>photoionization of a weak acid produces an ion which dissociates quite completely into a H^+ ion and the radical $H_{n-1}A$</u>. This result may be compared with the frequently observed enhancement of the strength of a weak acid upon excitation to its first electronic excited state [8].

Equation (5) is readily generalized (fig 1). The difference of reorganization free energies $\Delta^2(\Delta G_r)$ pertains to the photoionization of $H_{n-2}A^{2-}$ and $H_{n-1}A^-$, respectively. Since these two ions have comparable sizes, one expects from the Born equation that the ratio of ΔG_r's be approximately equal to the square of the ratio of ionic charges in the absence of dielectric saturation. The Born equation is approximate and there is saturation [3,6], especially for divalent ions, but one can safely conclude that $\Delta^2(\Delta G_r)$ is negative and not negligible. Hence, the condition $\Delta^2 - \Delta^2(\Delta G_r) > 0$ is satisfied for all the acids of table 1. The radical $H_{n-1}A$ produced by photoionization of $H_{n-1}A^-$ ion is unstable and the ion $H_{n-2}A^-$ resulting from photoionization of $H_{n-2}A^{2-}$ does not protonate to any extent. The same analysis and conclusion are applicable to photoelectron emission by the ions HA^{2-} and A^{3-} of table 1.

2. Weak bases and their cations

Weak bases in aqueous solution have lower threshold energies than their cations (table 1). One has (cf. eq (3)) for the base B and its cation BH^+

$$\Delta G_e(B) - \Delta G_e(BH^+) = [\Delta G(BH^+) - \Delta G(B)]$$
$$+ [\Delta G(B^+) - \Delta G(BH^{2+})] + [\Delta G_r(BH^{2+}) - \Delta G_r(B^+)], \qquad (7)$$

where $\Delta G(X)$ represents the free energy of formation of X in aqueous solution. The first term (< 0) on the right hand side of eq (7) is (in eV)

$$\Delta G_K^1 = \Delta G(BH^+) - \Delta G(B)$$
$$= -0.0592 \log K_p \quad , \qquad (8)$$

where K_p is the equilibrium constant for the protonation reaction $B + H^+ = BH^+$. The difference between the reorganization free energies $\Delta^1(\Delta G_r)$ is negative and not negligible because of the difference in ionic charge (and despite dielectric saturation).

One has

$$\iota^1 = E_t^1 - E_t^0 + \Delta G_K^1$$
$$= [\Delta G(BH^{2+}) - \Delta G(B^+)] - \iota^1(\Delta G_r),\tag{9}$$

where the superscripts o and 1 refer to B and BH^+, respectively, and the first term between brackets on the right hand side is the change of free energy for the protonation reaction, $B^+ + H^+ = BH^{2+}$. This change of free energy, equal to $\iota^1 + \iota^1(\Delta G_r)$, is negative for triethylamine and ethylenediamine since $\iota^1 \leq 0.2$ eV for these bases and $\iota^1(\Delta G_r) \leq -0.5$ eV (see above). Thus, the ion B^+ produced by photoionization of B protonates and the cation BH^{2+} is stable provided the pH is not too high. A similar condition is reached for the photoionization of the ion BH_2^+ of ethylenediamine (table 1). The quantity $\iota^1 + \iota^1(\Delta G_r)$ for aniline is probably positive or close to zero since $\iota^1 = 0.8$ eV and protonation of B^+ must be minor even in strongly acidic solution. Thus, photoionization of a weak base B (aniline) produces an ion B^+ which is a weaker base than B. A similar effect is frequently observed as a result of the excitation of a weak base to its first electronic excited state [8].

3. Weak acids as electron acceptors in electron transfer reactions

The emission yield Y of a solution is proportional to the emitter concentration in the absence of complications [1-5]. This proportionality relationship also holds for the squared slope S^2 of the linear extrapolation plot Y vs. $E^{0.5}$ (sec 1, n = 0.5 here). The plots of the squared slope S^2 against acid concentration for acetic or propionic acid actually exhibit a maximum (fig 2). This result, as was recently shown [9], is typical of electron transfer between the emitting species and an electron acceptor upon absorption of a photon. The emitter and electron acceptor are different molecules of the same acid in the present case. Thus, some electrons which

would be emitted into the vapor phase in the absence of electron transfer in solution are not emitted and the yield is lowered as a result of electron transfer.

The emission yield with electron transfer in solution was shown [9] to obey the following equation to a good approximation

$$Y = kC_m \exp\left(-C_a/C_c\right). \tag{10}$$

There k is a proportionality constant; C_m and C_a are the emitter and acceptor concentration, respectively; and C_c is a concentration which is characteristic of the kinetics of electron transfer. Dissociation of the acids of fig 2 can be neglected, and one has $C_m = C_a = C$ to a good approximation, where C is the analytical acid concentration. The plot of Y (or S^2) vs. C exhibits a maximum at $C = C_c$ according to eq (10), and the plot of log (S^2/C) vs. C is linear. These predictions are verified in fig 2. The concentration $C = C_c$ at the maximum of the S^2 vs. C plot defines a volume $1/C_c$ in which electron transfer is supposed to occur according to the model leading to eq (9) [9]. One has ca. $v = 0.4$ nm^3 for the data of fig 2, that is, a radius of ca. 0.5 nm if the volume v is supposed to be spherical. This length can be interpreted as the separation of two acid molecules undergoing electron transfer. Caution is in order since a simplified model was assumed in the derivation of eq (10), and the quantity v can be viewed as the product of an actual volume and a probability of electron transfer [9].

Emitter and acceptor need not be the same acid. Thus, the yield of acetic acid is lowered by addition of formic acid. The latter does not contribute to emission since it has a higher threshold energy (10.0 eV, table 1) than acetic acid (9.00 eV). Log S^2 at constant (2 M) acetic acid concentration indeed was found to decrease linearly with formic acid concentration from 0 to 7 M.

Aliphatic acids also lower the emission yield of inorganic anions (fig 3) and cations (fig 4). The slope S to the power 1/n is proportional to the emission yield, where n is the exponent in the extrapolation plot Y^n vs. E of the anions or cations. Thus, the plots of log $S^{1/n}$ vs. acceptor concentration at constant emitter concentration in fig 3 and 4 are linear as expected from eq (10). A strong anion effect on the rate of electron transfer is displayed in fig 3 just as in a similar plot in [9] for emission by halides in presence of Ba^{2+} ions (electron acceptor).

The results of fig 4 for Tl^+ ion are the first ones obtained for the lowering of the emission yield of a cation. The Tl^+ ion was selected because complexation with acetic or formic acid is negligible ($K_1 = 10^{-0.11}$ for acetic acid [10]). Lowering of the emission yield of anions by electron transfer to cations is common [9], e.g., the effect of Fe^{2+} on emission by Cl^-. The Fe^{2+} ions, however, do not affect emission by other Fe^{2+} ions because of electrostatic repulsion between Fe^{2+} ions (ionic atmosphere). Lowering of the yield of a cation (Tl^+) is observed with an electrically neutral electron acceptor (fig 4). Conversely, the yield of Tl^+ is not lowered by H^+ ion even in 6 M solution of a strong acid (trifluoroacetic), although H^+ ion is an excellent electron acceptor in the emission by anions.

Acknowledgment

References

[1] I. Watanabe, J. B. Flanagan and P. Delahay, J. Chem. Phys. 73 (1980) 2057.

[2] K. von Burg and P. Delahay, Chem. Phys. Letters 78 (1981) 287.

[3] P. Delahay, K. von Burg and A. Dziedzic, Chem. Phys. Letters 79 (1981) 157.

[4] P. Delahay and K. von Burg, Chem. Phys. Letters, submitted.

[5] A. M. Brodsky and A. V. Tsarevsky, J. Chem. Soc. Faraday Trans. II 72 (1976) 1781.

[6] R. M. Noyes, J. Am. Chem. Soc. 86 (1964) 971.

[7] B. Case and R. Parsons, Trans. Faraday Soc. 63 (1967) 1224.

[8] R. P. Bell, The proton in chemistry, 2nd ed. (Chapman and Hall, London, 1973, pp. 106-110).

[9] K. von Burg and P. Delahay, Chem. Phys. Letters, submitted.

[10] J. Bjerrum, G. Schwarzenbach and L. G. Sillen, Stability constants, part I, organic ligands (The Chemical Society, London, 1957, p. 3).

Table 1

Threshold energies and relevant quantities for weak acids and bases and their
ions[a]

Acid	E_t^j (eV)	ΔG_K^j (eV)	ϵ^j (eV)
formic	10.0 , 7.55	0.22	2.2
acetic	9.00, 7.82	0.28	0.9
monochloroacetic	9.24, 8.50	0.17	0.6
trichloroacetic	9.46, 8.55	0.04	0.9
propionic	9.08, 8.42	0.29	0.4
n-butyric	8.99, 8.23	0.29	0.5
carbonic	− , 9.07, 7.40	− , 0.61	− , 1.1
oxalic	8.26, 7.50, 7.32	0.08, 0.25	0.7, −0.1
tartaric	8.55, 7.72, 7.37	0.18, 0.26	0.6, 0.1
malonic	8.74, 8.33, 7.80	0.17, 0.34	0.2, 0.2
maleic	8.20, 8.31, 8.40	0.11, 0.36	−0.2, −0.4
phosphoric	9.45, 9.23, 8.79, 7.44	0.13, 0.43, 0.73	0.1, 0.0, 0.6
arsenic	9.44, 9.09, 8.50, 8.30	0.13, 0.40, 0.69	0.2, 0.2, −0.5
citric	8.66, 8.52, 8.39, 7.48	0.19, 0.28, 0.38	0.0, −0.1, 0.5
triethylamine	6.73, 7.57	−0.64	0.2
aniline	7.39, 8.44	−0.27	0.8
ethylenediamine	7.20, 7.47, 8.13	−0.59, −0.41	−0.3, 0.3

[a] 1 M solutions in all cases except for formic (4 M) and oxalic (0.4 M)
acids, oxalate (0.4 M), bitartrate (0.05 M), mono- and dibasic arsenate
(0.5 M), tribasic arsenate (0.25 M), aniline (0.05 M).

Captions to Figures

Fig. 1. Free energy diagram for photoelectron emission by H_nA, $H_{n-1}A^-$ and $H_{n-2}A^{2-}$. All species shown are in aqueous solution. See text for notations.

Fig. 2. Squared slope $S^2 = (dY^{0.5}/dE)^2$ for emission by acetic (A) and propionic (B) acid vs. acid concentration C. Points are experimental, and the curves were calculated from eq (10). Log (S^2/C) vs. acid concentration for acetic (C) and propionic (D) acids.

Fig. 3. Log $S^{1/n}$ for emission by anions vs. formic acid concentration for 2 M KCl (A), 2 M KBr (B), 2 M KI (C) and 1 M Li_2SO_4 (D). $S = dY^n/dE$ with n = 0.5 for Cl^-, SO_4^{2-} and n = 0.4 for Br^- and I^-.

Fig. 4. Log $S^{2.5}$ vs. acid concentration for emission by Tl^+ ions (0.25 M $TlClO_4$) in presence of trifluoroacetic (A), formic (B) or acetic (C) acid. $S = dY^{0.4}/dE$ for Tl^+ ion.

FIG. 1

FIG. 2

FIG. 3

FIG. 4

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1

Commander, Naval Air Systems Command
Attn: Code 310C (H. Rosenwasser)
Department of the Navy
Washington, D.C. 20360
	1	Naval Ship Research and Development
		   Center
		Attn: Dr. G. Bosmajian, Applied
		   Chemistry Division
		Annapolis, Maryland 21401
			1

Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco   96503   1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland   21402   1

472:GAN:716:ddc
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, 359

	No. Copies		No. Copies
Dr. Paul Delahav ~~Department of Chemistry New York University New York, New York 10003~~	1	Dr. P. J. Hendra Department of Chemistry University of Southhampton Southhampton SO9 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auborn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1		
Dr. T. Katan Lockheed Missiles & Space Co, Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1	Dr. R. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	No. Copies		No. Copies
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
S. Ruby DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton S09 5NH England	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	No. Copies		No. Copies
Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. John Kincaid Department of the Navy Stategic Systems Project Office Room 901 Washington, DC 20376	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	M. L. Robertson Manager, Electrochemical Power Sonices Division Naval Weapons Support Center	
Dr. M. G. Sceats Department of Chemistry University of Rochester Rochester, New York 14627	1	Crane, Indiana 47522	1
		Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory	
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	University of California Berkeley, California 94720	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University		Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709	1
Evanston, Illinois 60201	1	Dr. Denton Elliott Air Force Office of	
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1	Scientific Research Bldg. 104 Bolling AFB Washington, DC 20332	1
Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	1		

TECHNICAL REPORT DISTRIBUTION LIST, 051B

	No. Copies		No. Copies

Professor K. Wilson
Department of Chemistry,
  B-014
University of California,
  San Diego
La Jolla, California 92093
	1	Dr. B. Vonnegut
		State University of New York
		Earth Sciences Building
		1400 Washington Avenue
		Albany, New York 12203
			1

Professor C. A. Angell Department of Chemistry Purdue University West Lafayette, Indiana 47907	1	Dr. Hank Loos Laguna Research Laboratory 21421 Stans Lane Laguna Beach, California 92651	1
Professor P. Meijer Department of Physics Catholic University of America Washington, D.C. 20064	1	Dr. John Latham University of Manchester Institute of Science & Technology P.O. Box 88 Manchester, England M601QD	1
Dr. S. Greer Chemistry Department University of Maryland College Park, Maryland 20742	1		
Professor P. Delahay New York University 100 Washington Square East New York, New York 10003	1		
Dr. T. Ashworth Department of Physics South Dakota School of Mines & Technology Rapid City, South Dakota 57701	1		
Dr. G. Gross New Mexico Institute of Mining & Technology Socorro, New Mexico 87801	1		
Dr. J. Kassner Space Science Research Center University of Missouri - Rolla Rolla, Missouri 65401	1		
Dr. J. Telford University of Nevada System Desert Research Institute Lab of Atmospheric Physics Reno, Nevada 89507	1		